

## IMAGES IN CLINICAL PRACTICE

### MULTIPLE CONGENITAL ABNORMALITIES

Shweta Mishra, Piyush Chandel, Supriya Bisht

Department of Neonatology, Fortis Hospital, Noida, Uttar Pradesh, India

#### ARTICLE HISTORY

Received 4 May 2018

Accepted 5 September 2018

A 7 days old male infant was referred to us with sepsis, cardiovascular shock, and multiple congenital abnormalities. His birth weight was 2.24 kg and APGAR score was not known. At admission to our unit, he was very sick, had severe respiratory distress, shock (blood pressure was 30/18 mm of Hg), hypoxic and had active bleeding from the nasogastric tube. The abdomen was distended with decreased bowel sounds and there was hepatosplenomegaly. On cardiovascular system examination, he had a systolic murmur. He had dysmorphism in form of micrognathia, flat nasal bridge, high arched palate, congenital talipes equinovarus deformity, polydactyly (bifid right thumb), limb length discrepancy and bilateral undescended testis (Figure 1). He was put on mechanical ventilation. Inotropes, intravenous (IV) fluids and IV antibiotics were started. Echocardiography showed patent ductus arteriosus (PDA) with an atrial septal defect (ASD). Ultrasound (USG) abdomen and scrotum revealed bilateral undescended testis with right-sided hydrocele. USG cranium revealed hypoplasia of the corpus callosum. Blood culture grew *Enterobacter cloacae*. Coagulation profile was abnormal. He was given multiple platelets, fresh frozen plasma and packed red blood cell transfusions. Gradually he improved and was extubated on day 4 of admission. A repeat echocardiography on Day 9 of admission showed closure of PDA and only presence of ASD 5.5 mm. After stabilization, MRI brain was done that showed lobar holoprosencephaly, hypoplastic corpus callosum, pons and cerebellum with the prominent fourth ventricle. TORCH titers were negative and ophthalmological evaluation was normal. Karyotype analysis proved the diagnosis.

*What is the syndrome?*



**CONTACT** Dr. Shweta Mishra

**Email:** [shwetapaed@gmail.com](mailto:shwetapaed@gmail.com)

**Address for Correspondence:** Dr. Shweta Mishra, 8/101, Shakti Kunj Apartment, B9/3, Sec 62, Noida, Uttar Pradesh, India.

©2019 Pediatric Oncall

Karyotype analysis of the patient revealed 46 XY, del (13q22 -32). It is also known as Orbeli syndrome based on the Russian physician who described the syndrome in 1971. It is a partial monosomy involving the long arm of chromosome 13. Patient presents with varying phenotypic features as per deleted band of the chromosome 13q and are classified into 3 groups. Deletions involving chromosome 13 proximal to q32 band can result in phenotypes such as mild mental retardation, growth delay and various dysmorphic features. When loci of q32 gets deleted, it leads to severe congenital malformations including brain anomalies. teters form is deletion distal to q32 which leads to severe mental retardation, but no brain malformation or growth delay.<sup>1,2</sup> The ZIC2 (zinc finger protein of cerebellum 2) gene has been mapped in this region and absence of it by deletion is considered to lead to brain anomaly like holoprosencephaly.

Various clinical features of this syndrome are low birth weight, growth failure, psychomotor retardation, craniofacial abnormalities, microcephaly, brachycephaly, hypertelorism, upslanting palpebral fissures, high arched palate, coloboma, prominent nasal bridge, clinodactyly, syndactyly, dislocation of hips, a , congenital heart disease, renal abnormalities, cryptorchidism, hypospadias and anal atresia.<sup>2</sup> There is no specific treatment was given exact an is unknown. Children with 13q deletion syndrome should be followed up continuously by a neurologist, pediatric surgeon, and a rehabilitation physician, who can also coordinate interventions with different specialists.

#### Compliance with Ethical Standards

**Funding:** None

**Conflict of Interest:** None

#### References :

1. Brown S, Gersen S, Anyane-Yeboah K, Warburton D. Preliminary definition of a "critical region" of chromosome 13 in q32: report of 14 cases with 13q deletions and review of the literature. *Am J Med Genet.* 1993; 45: 52-9.
2. Brown S, Russo J, Chitayat D, Warburton D. The 13q-syndrome: the definition of a critical deletion region in band 13q32. *Am J Hum Genet.* 1995; 57: 859-66