

## LETTER TO EDITOR (VIEWERS CHOICE)

**SUCCESSFUL SURGICAL REPAIR OF SUB-CORONAL HYPOSPADIAS IN A 10-YEAR-OLD MALE: A CASE OF DELAYED PRESENTATION AND EFFECTIVE INTERVENTION**Shashi Prakash<sup>1</sup>, Archana Yadav<sup>2</sup>.<sup>1</sup>Nursing Faculty, Govt. College of Nursing, S. N. Medical College, Agra, Uttar Pradesh, India,<sup>2</sup>B.Sc. Nursing Student, Govt. College of Nursing, S. N. Medical College, Agra, Uttar Pradesh, India.**KEYWORDS**

Sub-coronal hypospadias, TIP urethroplasty, Hypospadias repair, Pediatric urology, Surgical outcomes, Delayed presentation.

**ARTICLE HISTORY**

Received 12 Dec 2024

Accepted 03 Jan 2025

In order to demonstrate the difficulties and positive results of surgical restoration in a delayed presenting situation, we offer a case of sub-coronal hypopadias in a male patient aged ten. One of the most prevalent congenital abnormalities affecting male genitalia is hypospadias, which is usually identified and surgically fixed in early childhood. However, in settings with limited resources, delayed diagnosis and treatment are frequently seen, which raises serious social, psychological, and functional issues. In our instance, a tubularized incised plate (TIP) urethroplasty was performed on a male infant who had an aberrant urine stream and ventral curvature, with positive functional and cosmetic outcomes.

The urethral meatus's aberrant placement, which can vary from the glans to the perineum, is what defines hypopadias. Although the condition's occurrence is roughly 1 in 200 live male births, the way it manifests itself varies greatly depending on the location. Early diagnosis enables successful corrective surgery to be carried out in infancy in wealthy nations. However, delayed diagnoses are typical in nations like India where issues like poverty, illiteracy, and restricted access to healthcare continue to exist. A later presentation in these situations, usually in childhood or even adulthood, may result in more issues.<sup>1</sup>

The subject of this study is a 10-year-old boy who has experienced urinating issues since he was a little toddler. His symptoms included spraying, an erratic urine stream, and the need to crouch when urinating because of his penis' ventral curvature. Due to logistical and familial limitations, the illness was not treated during infancy despite these symptoms. By the time the youngster was ten years old, the illness had caused him considerable distress, particularly after he had been teased by classmates at school, which made his social anxiety much worse.

No other severe health issues were found, even though the patient had a history of cryptorchidism that required

orchiopexy surgery at 17 months. A hooded prepuce and a sub-coronal meatus with a significant chordee were discovered during the patient's examination. His blood and urine tests were normal, and ultrasonography confirmed the diagnosis of sub-coronal hypospadias without any concomitant abnormalities.

The TIP urethroplasty technique, which is known for its exceptional results in treating hypopadias, was used to plan the surgical intervention.<sup>2</sup> The urethral plate was mobilized, the chordee was corrected by dorsal plication, and the plate was tubularized to create a new urethra. Glanuloplasty was performed, and local skin flaps were used to cover the area. The patient was kept on postoperative monitoring for potential complications, such as bleeding, fistula formation, or infection, and a urethral catheter was placed for seven days to promote healing.

The outcomes of the six-month follow-up were outstanding. The patient had a straight penis, a normal urine stream, and a visually appealing outcome. There were no signs of stricture, chordee recurrence, or fistula. Preoperative counseling also addressed the psychological effects of the disease and after surgery, the patient's self-esteem significantly improved and he no longer experienced any social challenges associated with his condition.

This case illustrates several important aspects of hypospadias management. First, it highlights the issue of delayed presentation, which remains common in resource-limited regions. The impact on a child's emotional and social well-being can be significant, as was evident in this case. Second, it highlights how well the TIP urethroplasty technique - which has emerged as the gold standard for treating sub-coronal hypospadias-works, even in older kids.<sup>3</sup> In these situations, the method is the best option due to its ease of use, low rate of complications, and superior functional and cosmetic results.<sup>4</sup> Lastly, the story emphasizes how critical it is to address the social and psychological difficulties that children with congenital conditions like hypopadias experience. A multidisciplinary approach, psychological assistance, and preoperative counseling are essential for guaranteeing the patient's general wellbeing.

In conclusion, this case shows that surgical treatment of sub-coronal hypospadias can produce outstanding

**Address for Correspondance:** Shashi Prakash, Block-A, Govt. College of Nursing S. N. Medical College, Agra, Uttar Pradesh, India-282002.

**Email:** prakashshashipr@gmail.com

©2025 Pediatric Oncall

functional and cosmetic results, even if presentation is delayed. For older children, the TIP urethroplasty technique is still a dependable alternative. Effective therapy requires prompt intervention as well as consideration of the psychological aspects of the illness.<sup>5</sup>

**Compliance with Ethical Standards**

**Funding** None

**Conflict of Interest** None

**References:**

1. Snodgrass W. Tubularized, incised plate urethroplasty for distal hypospadias. *J Urol.* 1994 Feb;151(2):464-5. doi: 10.1016/s0022-5347(17)34991-1. PMID: 8283561.
2. Yang L, Zhang Q, Zhang Z, et al. Meta-analysis comparing TIP urethroplasty and other techniques for hypospadias repair. *J Urol.* 2018;199(4):986-993.
3. Zheng C, Liu L, Yang X, et al. Comparison of TIP and Onlay Island Flap for hypospadias repair: A systematic review and meta-analysis. *Urology.* 2017;108:17-23.
4. Duckett JW, Wilcox DT. The tubularized island flap in the repair of hypospadias. *J Urol.* 1996;156(1):224-229.
5. Sharma R, Li S, Singh P, et al. Stem cell therapy in hypospadias repair: Potential future directions. *Urol Ann.* 2020;12(1):7-12.